



Contribution ID: 32

Type: **contributed parallel talk**

Three-Body Unitary Coupled-Channel Analysis on $\eta(1405/1475)$

Thursday 20 July 2023 15:40 (20 minutes)

The recent BESIII data on $J/\psi \rightarrow \gamma(K_S K_S \pi^0)$, which is significantly more precise than earlier $\eta(1405/1475)$ -related data, enables quantitative discussions on $\eta(1405/1475)$ at the previously unreachable level. We conduct a three-body unitary coupled-channel analysis of experimental Monte-Carlo outputs for radiative J/ψ decays via $\eta(1405/1475): K_S K_S \pi^0$ Dalitz plot distributions from the BESIII, and branching ratios of $\gamma(\eta\pi^+\pi^-)$ and $\gamma(\eta\pi^+\pi^-)$ final states relative to that of $\gamma(K\bar{K}\pi)$. Our model systematically considers (multi-)loop diagrams and an associated triangle singularity, which is critical in making excellent predictions on $\eta(1405/1475) \rightarrow \pi\pi\pi$ lineshapes and branching ratios. The $\eta(1405/1475)$ pole locations are revealed for the first time. Two poles for $\eta(1405)$ are found on different Riemann sheets of the $K^*\bar{K}$ channel, while one pole for $\eta(1475)$. The $\eta(1405/1475)$ states are described with two bare states dressed by continuum states. The lower bare state would be an excited η' , while the higher one could be an excited $\eta^{(1)}$, hybrid, glueball, or their mixture. This work presents the first-ever pole determination based on a manifestly three-body unitary coupled-channel framework applied to experimental three-body final state distributions (Dalitz plots). This presentation is based on arXiv:2212.07904 to appear in Phys. Rev. D.

Consent

I consent to recording/broadcasting my presentation.

Primary author: NAKAMURA, Satoshi (Shandong University)

Presenter: NAKAMURA, Satoshi (Shandong University)

Session Classification: Parallel B

Track Classification: spectroscopy