



Contribution ID: 45

Type: **contributed parallel talk**

Precise determination of the decay rates of $\eta_c \rightarrow \gamma\gamma$, $J/\psi \rightarrow \gamma\eta_c$ and $J/\psi \rightarrow \eta_c e^+ e^-$ from lattice QCD

Tuesday 18 July 2023 14:20 (20 minutes)

We present results from our calculation of decays rates for $\eta_c \rightarrow \gamma\gamma$, $J/\psi \rightarrow \gamma\eta_c$ and $J/\psi \rightarrow \eta_c e^+ e^-$ in lattice QCD with the effect of u, d, s and c quarks in the sea for the first time. We use the Highly Improved Staggered Quark formulism, four values of the lattice spacing and sea u/d quarks down to their physical values. Our results are accurate at the 1-2% level and are therefore now more accurate than results from experiment. We find $\Gamma(\eta_c \rightarrow \gamma\gamma) = 6.788(45)_{\text{fit}}(41)_{\text{syst}}$ keV, which agrees well with experimental results for $\gamma\gamma \rightarrow \eta_c \rightarrow K\bar{K}\pi$. This is in tension with the global PDG fit at the 4σ level, however, and we therefore advise this fit is revisited. We find $\Gamma(J/\psi \rightarrow \gamma\eta_c) = 2.219(17)_{\text{fit}}(18)_{\text{syst}}(24)_{\text{expt}}(4)_{\text{QED}}$ keV, which agrees well with results from CLEO. Finally, we predict $\Gamma(J/\psi \rightarrow \eta_c e^+ e^-) = 0.01349(21)_{\text{latt}}(13)_{\text{QED}}$ keV. We compare our results with other theoretical approaches, while simple relationships between form factors and the J/ψ decay constant in the nonrelativistic limit are also tested.

Consent

I consent to recording/broadcasting my presentation.

Primary authors: COLQUHOUN, Brian (University of Glasgow); COOPER, Laurence (University of Glasgow); DAVIES, Christine (University of Glasgow); LEPAGE, G. Peter (Cornell University)

Presenter: COLQUHOUN, Brian (University of Glasgow)

Session Classification: Parallel A

Track Classification: decays