

Quark-Hadron Duality Violation and Higher Order $1/m_b$ corrections in inclusive $B \rightarrow X_c \ell \bar{\nu}$

Quirks in Quark Flavour Physics 2024 - Zadar, Croatia

Ilija S. Milutin¹

in collaboration with
Thomas Mannel¹, Rens Verkade² and K. Keri Vos²


¹CPPS, Universität Siegen, Germany

²GWFP, Maastricht University and Nikhef, The Netherlands



The V_{cb} puzzle

- Inclusive $B \rightarrow X_c \ell \bar{\nu}$ for extraction of V_{cb}
→ See talk by Keri Vos
- Heavy Quark Expansion (HQE) → power series in Λ_{QCD}/m_b
Chay, Georgi, Bigi, Shifman, Uraltsev, Vainstain, Manohar, Wise, Neubert, Mannel,...
- Split b quark momentum as $p_b = m_b v + k$
→ Expand in $k \sim iD$



Bernlochner, Prim, Vos (*Eur. Phys. J. Spec. Top.* (2024))

Matrix elements

- Perform Operator Product Expansion (OPE)

Chay, Georgi, Bigi, Shifman, Uraltsev, Vainstain, Manohar, Wise, Neubert, Mannel,...

$$d\Gamma = d\Gamma^{(3)} + \frac{1}{m_b^2} d\Gamma^{(5)} + \frac{1}{m_b^3} d\Gamma^{(6)} + \frac{1}{m_b^4} d\Gamma^{(7)} + \dots, \quad d\Gamma^{(n)} = \sum_i C_i^{(n)} \langle B | \mathcal{O}_i^{(n)} | B \rangle$$

- $d\Gamma^{(3)}$: Partonic result ($d\Gamma^{(4)} = 0$ due to Heavy Quark Symmetries)
- $d\Gamma^{(5)}$: 2 parameters

$$2m_B \mu_{\pi}^2 = -\langle B | \bar{b}_v (iD)^2 b_v | B \rangle$$

$$2m_B \mu_G^2 = \langle B | \bar{b}_v (-i\sigma^{\mu\nu}) (iD_\mu) (iD_\nu) b_v | B \rangle$$

- $d\Gamma^{(6)}$: 2 parameters


$$2m_B \rho_D^3 = \langle B | \bar{b}_v [iD_\mu, [ivD, iD^\mu]] b_v | B \rangle / 2$$

$$2m_B \rho_{LS}^3 = \langle B | \bar{b}_v [iD_\mu, [ivD, iD_\nu]] (-i\sigma^{\mu\nu}) b_v | B \rangle / 2$$

- $d\Gamma^{(7)}$: 9 parameters (at tree level) Mannel, Turczyk, Uraltsev [1009.4622]; Kobach, Pal [1704.00008]
- $d\Gamma^{(8)}$: 18 parameters (at tree level) Mannel, Turczyk, Uraltsev [1009.4622]; Kobach, Pal [1704.00008]

Counting Operators with Reparametrization Invariance

- In HQE, choice of v_μ is not unique
Dugan, Golden, Grinstein, Chen, Luke, Manohar, Hill, Solon, Heinonen,...
- Lorentz invariance of QCD →
Reparametrization Invariance (RPI)
imposed by $v_\mu \rightarrow v_\mu + \delta v_\mu$
- **RPI relates different orders** in $1/m_b$ expansion
Mannel, Vos [1802.09409]
→ This allows us to find combinations of operators which are RPI
- Up to $1/m_b^4$: total of 8 independent parameters
Mannel, Vos [1802.09409]
- At $1/m_b^5$, we find **only 10 RPI operators**
Mannel, ISM, Vos [2311.12002]



Inclusive V_{cb} determination

- Want an RPI-observable \rightarrow only depend on reduced set of RPI operators

Inclusive V_{cb} determination

- Want an RPI-observable \rightarrow only depend on reduced set of RPI operators
- Dilepton invariant mass (q^2) moments** are RPI and can be used to extract $|V_{cb}^{\text{incl}}|$
(q^2 -cut needed due to experimental setup)

Bernlochner, Welsch, Fael, Olschewsky, Persson, von Tonder, Vos [2205.10274]

$$\langle (q^2)^n \rangle_{\text{cut}} = \frac{\int_{q^2 > q_{\text{cut}}^2} dq^2 (q^2)^n \frac{d\Gamma}{dq^2}}{\int_{q^2 > q_{\text{cut}}^2} dq^2 \frac{d\Gamma}{dq^2}}$$

Inclusive V_{cb} determination

- Want an RPI-observable \rightarrow only depend on reduced set of RPI operators
- Dilepton invariant mass (q^2) moments** are RPI and can be used to extract $|V_{cb}^{\text{incl}}|$
(q^2 -cut needed due to experimental setup)

Bernlochner, Welsch, Fael, Olschewsky, Persson, von Tonder, Vos [2205.10274]

$$\langle (q^2)^n \rangle_{\text{cut}} = \frac{\int_{q^2 > q_{\text{cut}}^2} dq^2 (q^2)^n \frac{d\Gamma}{dq^2}}{\int_{q^2 > q_{\text{cut}}^2} dq^2 \frac{d\Gamma}{dq^2}}$$

- Data \rightarrow values for reduced set of RPI parameters up to $1/m_b^4 \rightarrow Br(\bar{B} \rightarrow X_c \ell \bar{\nu}) \rightarrow |V_{cb}^{\text{incl}}| = (41.69 \pm 0.63) \times 10^{-3}$

Bernlochner, Vos, et al. [2205.10274]

Inclusive V_{cb} determination

- Want an RPI-observable \rightarrow only depend on reduced set of RPI operators
- Dilepton invariant mass (q^2) moments** are RPI and can be used to extract $|V_{cb}^{\text{incl}}|$
(q^2 -cut needed due to experimental setup)

Bernlochner, Welsch, Fael, Olschewsky, Persson, von Tonder, Vos [2205.10274]

$$\langle (q^2)^n \rangle_{\text{cut}} = \frac{\int_{q^2 > q_{\text{cut}}^2} dq^2 (q^2)^n \frac{d\Gamma}{dq^2}}{\int_{q^2 > q_{\text{cut}}^2} dq^2 \frac{d\Gamma}{dq^2}}$$

- Data \rightarrow values for reduced set of RPI parameters up to $1/m_b^4 \rightarrow Br(\bar{B} \rightarrow X_c \ell \bar{\nu}) \rightarrow |V_{cb}^{\text{incl}}| = (41.69 \pm 0.63) \times 10^{-3}$

Bernlochner, Vos, et al. [2205.10274]

- First determination** of V_{cb} up to $\mathcal{O}(1/m_b^4)$ and **first extraction** of $1/m_b^4$ matrix elements from data
- Agreement at $1 - 2\sigma$ level with previous $\mathcal{O}(1/m_b^3)$ determinations

Finauri, Gambino [2310.20324]; Bordone, Capdevila, Gambino [2107.00604]; Alberti, Gambino, Healey, Nandi[1411.6560]; Gambino, Schwanda [1307.4551]

Where do we currently stand?

Γ	tree	α_s	α_s^2	α_s^3
Partonic	✓	✓	✓	✓
$1/m_b^2$	✓	✓		
$1/m_b^3$	✓	✓		
$1/m_b^4$	✓			
$m_b^{\text{kin}}/\bar{m}_c$		✓	✓	✓

$\langle (q^2)^n \rangle$	tree	α_s	α_s^2	α_s^3
Partonic	✓	✓		
$1/m_b^2$	✓	✓		
$1/m_b^3$	✓	✓		
$1/m_b^4$	✓			

- **Green:** known perturbative corrections Jezabek, Kuhn (1989); Melnikov (2008); Pak, Czarnecki (2008); Becher, Boos, Lunghi (2007); Alberti, Gambino, Nandi (2014); Mannel, Pivovarov, Rosenthal (2015); Gambino, Healey, Turczyk (2016); Mannel, Pivovarov (2020); Fael, Schonwald, Steinhauser (2020, 2021)
- Next step for higher precision:
 - $1/m_b^5$
 - Quark-Hadron Duality Violation

Going higher in the $1/m_b$ expansion

- Dimension 8 contains: Bigi, Mannel, Turczyk, Uraltsev [0911.3322]


“genuine”

$$\frac{1}{m_b^5}$$

&

Intrinsic Charm


$$\frac{1}{m_b^3 m_c^2}$$



Going higher in the $1/m_b$ expansion

- Dimension 8 contains: Bigi, Mannel, Turczyk, Uraltsev [0911.3322]

“genuine” $\frac{1}{m_b^5}$ & **Intrinsic Charm** $\frac{1}{m_b^3 m_c^2}$




- Numerically: $m_c^2 \sim m_b \Lambda_{\text{QCD}}$
- Dimension 8 contains enhanced terms which contribute like $1/m_b^4$ terms

Going higher in the $1/m_b$ expansion

- Dimension 8 contains: Bigi, Mannel, Turczyk, Uraltsev [0911.3322]


“genuine” $\frac{1}{m_b^5}$ & Intrinsic Charm $\frac{1}{m_b^3 m_c^2}$



- Numerically: $m_c^2 \sim m_b \Lambda_{\text{QCD}}$
- Dimension 8 contains enhanced terms which contribute like $1/m_b^4$ terms
- We need $1/m_b^3 \times 1/m_c^2$ contributions to complete the calculation at $1/m_b^4$
 - We calculate the full dimension 8 contributions
 - We extract the Intrinsic Charm contribution
 - We find that only 1 combination of parameters describes the IC in the q^2 -moments


Mannel, ISM, Vos [2311.12002]

q^2 -moments




Mannel, ISM, Vos [2311.12002]

q^2 -moments




Mannel, ISM, Vos [2311.12002]

q^2 -moments



Mannel, ISM, Vos [2311.12002]

q^2 -moments



Mannel, ISM, Vos [2311.12002]

Conclusions on dimension 8 contributions

- We identified the **10 RPI operators** at $1/m_b^5$

Conclusions on dimension 8 contributions

- We identified the **10 RPI operators** at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions

Conclusions on dimension 8 contributions

- We identified the **10 RPI operators** at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an **unexpectedly small** overall contribution of the dimension-8 operators

Conclusions on dimension 8 contributions

- We identified the 10 RPI operators at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an unexpectedly small overall contribution of the dimension-8 operators
- q^2 , E_ℓ , M_X^2 moments to $1/m_b^5$ available in open-source library `kolya`


Conclusions on dimension 8 contributions

- We identified the 10 RPI operators at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an unexpectedly small overall contribution of the dimension-8 operators
- q^2, E_ℓ, M_X^2 moments to $1/m_b^5$ available in open-source library `kolya`
- We provide so-called trace formulae for determinations of observables to $\mathcal{O}(1/m_b^5)$ for $B \rightarrow X_c \ell \bar{\nu}$ Mannel, ISM, Vos [2311.12002]


Conclusions on dimension 8 contributions

- We identified the 10 RPI operators at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an unexpectedly small overall contribution of the dimension-8 operators
- q^2 , E_ℓ , M_X^2 moments to $1/m_b^5$ available in open-source library `kolya`
- We provide so-called trace formulae for determinations of observables to $\mathcal{O}(1/m_b^5)$ for $B \rightarrow X_c \ell \bar{\nu}$ Mannel, ISM, Vos [2311.12002]
- Work in progress: Full $|V_{cb}|^{\text{incl}}$ fit to $1/m_b^5$


Quark-Hadron Duality (QHD)



Quark-Hadron Duality (QHD)




Quark-Hadron Duality (QHD)




- QHD in OPE: well behaved Taylor series and analytic in Λ_{QCD}/Q where $Q = m_b v - q$

Quark-Hadron Duality (QHD)



- QHD in OPE: well behaved Taylor series and analytic in Λ_{QCD}/Q where $Q = m_b v - q$
→ data currently shows no indications of failure of HQE

Quark-Hadron Duality (QHD)



- QHD in OPE: well behaved Taylor series and analytic in Λ_{QCD}/Q where $Q = m_b v - q$
 - data currently shows no indications of failure of HQE
 - but factorially increasing number of HQE parameters hint for **QHD Violation**

Modelling QHDV

- Optical theorem:

$$d\Gamma \propto L^{\mu\nu} \operatorname{Im} [T_{\mu\nu}(vQ, Q^2)]$$

- leptonic tensor $L^{\mu\nu}$
- hadronic tensor as imaginary part of time-ordered product:

$$T_{\mu\nu}(Q) = \int d^4x e^{-iQ \cdot x} \langle B(p) | T\{\bar{b}_\nu(x)\Gamma_\mu c(x) \bar{c}(0)\bar{\Gamma}_\nu b_\nu(0)\} | B(p) \rangle$$

Modelling QHDV

- Optical theorem:

$$d\Gamma \propto L^{\mu\nu} \operatorname{Im} [T_{\mu\nu}(vQ, Q^2)]$$

- leptonic tensor $L^{\mu\nu}$
- hadronic tensor as imaginary part of time-ordered product:

$$T_{\mu\nu}(Q) = \int d^4x e^{-iQ \cdot x} \langle B(p) | T\{\bar{b}_\nu(x)\Gamma_\mu c(x) \bar{c}(0)\bar{\Gamma}_\nu b_\nu(0)\} | B(p) \rangle$$

- Expand external field propagator for the charm quark:

$$\begin{aligned} -iS_{\text{BGF}} &= \frac{1}{\not{Q} + i\not{D} - m_c} \\ &= \frac{1}{\not{Q} - m_c} - \frac{1}{\not{Q} - m_c} (i\not{D}) \frac{1}{\not{Q} - m_c} + \dots \\ &= \sum_{k=0}^{\infty} \left(\frac{1}{Q^2}\right)^{k+1} \not{Q} \left[- (i\not{D}) \not{Q} \right]^k \quad (m_c = 0) \end{aligned}$$

Modelling QHDV continued

- Taking forward matrix element with B meson with velocity v :

$$T_{\mu\nu}(Q) = \sum_{k=0}^{\infty} \left(\frac{1}{Q^2} \right)^{k+1} \langle B(v) | \bar{b}_v \Gamma_\mu \not{Q} [-(i\not{D}) \not{Q}]^k \bar{\Gamma}_\nu b_v(0) | B(v) \rangle$$

Modelling QHDV continued

- Taking forward matrix element with B meson with velocity v :

$$T_{\mu\nu}(Q) = \sum_{k=0}^{\infty} \left(\frac{1}{Q^2} \right)^{k+1} \langle B(v) | \bar{b}_v \Gamma_\mu \not{Q} [-(i\not{D}) \not{Q}]^k \bar{\Gamma}_\nu b_v(0) | B(v) \rangle$$

- Order by order (schematically, all Lorentz indices suppressed):

$$\langle B(v) | \bar{b}_v \Gamma \not{Q} \bar{\Gamma} b_v | B(v) \rangle = a_0^{(i,0)}(vQ) ,$$

$$\langle B(v) | \bar{b}_v (-1) \Gamma \not{Q} (i\not{D}) \not{Q} \bar{\Gamma} b_v | B(v) \rangle = \Lambda_{\text{HQE}} \left(a_0^{(i,1)}(vQ)^2 + a_1^{(i,1)} Q^2 \right) ,$$

$$\langle B(v) | \bar{b}_v \Gamma \not{Q} (i\not{D}) \not{Q} (i\not{D}) \not{Q} \bar{\Gamma} b_v | B(v) \rangle = \Lambda_{\text{HQE}}^2 \left(a_0^{(i,2)}(vQ)^3 + a_1^{(i,2)}(vQ) Q^2 \right) ,$$

...

→ the index $i = 1, \dots, 5$ denotes the five scalar components of $T_{\mu\nu}$

Modelling QHDV continued

- Introduce dimensionless variables $r^2 = Q^2/\Lambda_{\text{HQE}}^2$ and $t = vQ/\Lambda_{\text{HQE}}$:

$$T_i(t, r^2) = \frac{1}{\Lambda_{\text{HQE}}} \sum_{l=0}^{\infty} \left(\frac{1}{r^2} \right)^{l+1} P_l^{(i)}(t), \quad P_l^{(i)}(t) = \sum_{k=0}^{l+1} t^{l+1-k} a_k^{(i, k+l)}$$

Modelling QHDV continued

- Introduce dimensionless variables $r^2 = Q^2/\Lambda_{\text{HQE}}^2$ and $t = vQ/\Lambda_{\text{HQE}}$:

$$T_i(t, r^2) = \frac{1}{\Lambda_{\text{HQE}}} \sum_{l=0}^{\infty} \left(\frac{1}{r^2} \right)^{l+1} P_l^{(i)}(t), \quad P_l^{(i)}(t) = \sum_{k=0}^{l+1} t^{l+1-k} a_k^{(i, k+l)}$$

- We can study coefficients $a_k^{(i, k+l)}$ using known contributions to $1/m_b^5$

Modelling QHDV continued

- Introduce dimensionless variables $r^2 = Q^2/\Lambda_{\text{HQE}}^2$ and $t = vQ/\Lambda_{\text{HQE}}$:

$$T_i(t, r^2) = \frac{1}{\Lambda_{\text{HQE}}} \sum_{l=0}^{\infty} \left(\frac{1}{r^2} \right)^{l+1} P_l^{(i)}(t), \quad P_l^{(i)}(t) = \sum_{k=0}^{l+1} t^{l+1-k} a_k^{(i, k+l)}$$

- We can study coefficients $a_k^{(i, k+l)}$ using known contributions to $1/m_b^5$
→ model factorial growth as $P_l(t) = (2l)! p_l(t)$

Modelling QHDV continued

- Introduce dimensionless variables $r^2 = Q^2/\Lambda_{\text{HQE}}^2$ and $t = vQ/\Lambda_{\text{HQE}}$:

$$T_i(t, r^2) = \frac{1}{\Lambda_{\text{HQE}}} \sum_{l=0}^{\infty} \left(\frac{1}{r^2} \right)^{l+1} P_l^{(i)}(t), \quad P_l^{(i)}(t) = \sum_{k=0}^{l+1} t^{l+1-k} a_k^{(i, k+l)}$$

- We can study coefficients $a_k^{(i, k+l)}$ using known contributions to $1/m_b^5$
 - model factorial growth as $P_l(t) = (2l)! p_l(t)$
 - use known contributions to make ansatz for polynomials $p_l(t)$

$$p_l^{(1,4)}(t) = t^{l+1} + t^l + \dots + t$$

$$p_l^{(2,3)}(t) = t^l + t^{l-1} + \dots + t + 1$$

$$p_l^{(5)}(t) = t^{l+1} + t^l + \dots + t^2$$

Borel Transform

- Now perform **Borel Transform** of $T_{\mu\nu}(r^2, t)$ w.r.t. $\lambda = 1/r$ and transform back
- Schematically:

$$F(\lambda) = \sum_n (2n)! \lambda^{2n}$$

$$B(M) = \sum_n M^{2n} = \frac{1}{1 - M^2} = \frac{1}{1 + M} \frac{1}{1 - M}$$

$$F(\lambda) = \int_0^\infty dM e^{-M} B(\lambda M)$$

Borel Transform

- Now perform **Borel Transform** of $T_{\mu\nu}(r^2, t)$ w.r.t. $\lambda = 1/r$ and transform back
- Schematically:

$$F(\lambda) = \sum_n (2n)! \lambda^{2n}$$

$$B(M) = \sum_n M^{2n} = \frac{1}{1 - M^2} = \frac{1}{1 + M} \frac{1}{1 - M}$$

$$F(\lambda) = \int_0^\infty dM e^{-M} B(\lambda M)$$

- Singularity creates ambiguity → **define duality violation** as difference between the two integration prescriptions (taking into account possible factors of t):

$$\frac{1}{1 - M + i\epsilon} - \frac{1}{1 - M - i\epsilon} = 2i\pi\delta(1 - M)$$

Borel Transform

- Now perform **Borel Transform** of $T_{\mu\nu}(r^2, t)$ w.r.t. $\lambda = 1/r$ and transform back
- Schematically:

$$F(\lambda) = \sum_n (2n)! \lambda^{2n}$$

$$B(M) = \sum_n M^{2n} = \frac{1}{1 - M^2} = \frac{1}{1 + M} \frac{1}{1 - M}$$

$$F(\lambda) = \int_0^\infty dM e^{-M} B(\lambda M)$$

- Singularity creates ambiguity → **define duality violation** as difference between the two integration prescriptions (taking into account possible factors of t):

$$\frac{1}{1 - M + i\epsilon} - \frac{1}{1 - M - i\epsilon} = 2i\pi\delta(1 - M)$$

- Finally, **analytically continue** to the Minkowskian case by $\lambda \rightarrow i\kappa = i\Lambda_{\text{HQE}}/\sqrt{Q^2}$



The QHDV model

- This will lead to: Mannel, ISM, Verkade, Vos [WIP]

$$-\frac{1}{\pi} \hat{\Delta}_{DV} \text{Im} [T_{1,4}(vQ, Q^2)] = \\ C_{DV} \frac{N}{\Lambda_{HQE} - vQ} \frac{vQ}{\sqrt{Q^2}} \left(\sin \left(\frac{\sqrt{Q^2}}{\Lambda_{HQE}} \right) - \sqrt{\frac{vQ}{\Lambda_{HQE}}} \sin \left(\frac{1}{\sqrt{\Lambda_{HQE}}} \sqrt{\frac{Q^2}{vQ}} \right) \right)$$

- similar expressions for $T_{2,3,5}$
- N is a normalisation and C_{DV} characterises “strength” of QHDV
- $\Lambda_{HQE}=0.5$ GeV as default (based on HQE parameters)

Differential rates for QHDV



- Γ_0 : partonic rate, $\hat{q}^2 = q^2/m_b^2$, $y = 2E_\ell/m_b$

Conclusions and outlook

- We identified the 10 RPI operators at $1/m_b^5$
 - The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
 - We find an unexpectedly small overall contribution of the dimension-8 operators
 - q^2 , E_ℓ , M_X^2 moments to $1/m_b^5$ available in open-source library `kolya`
 - We provide so-called trace formulae for determinations of observables to $\mathcal{O}(1/m_b^5)$ for $B \rightarrow X_c \ell \bar{\nu}$ Mannel, ISM, Vos [2311.12002]
-

Conclusions and outlook


- We identified the 10 RPI operators at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an unexpectedly small overall contribution of the dimension-8 operators
- q^2, E_ℓ, M_X^2 moments to $1/m_b^5$ available in open-source library `kolya`
- We provide so-called trace formulae for determinations of observables to $\mathcal{O}(1/m_b^5)$ for $B \rightarrow X_c \ell \bar{\nu}$
Mannel, ISM, Vos [2311.12002]
- We built a model for QHDV guided by known contributions to $1/m_b^5$
Mannel, ISM, Verkade, Vos [soon!]
 - We model the factorial growth of the number of HQE parameters
 - We make an educated ansatz for the vQ dependence of $T_{\mu\nu}$ based on known contributions
 - We define the QHDV using the ambiguity arising from the inverse Borel transform

Conclusions and outlook

- We identified the 10 RPI operators at $1/m_b^5$
- The $1/m_b^3 \times 1/m_c^2$ contributions are (partially) cancelled by the strict $1/m_b^5$ contributions
- We find an unexpectedly small overall contribution of the dimension-8 operators
- q^2, E_ℓ, M_X^2 moments to $1/m_b^5$ available in open-source library `kolya`
- We provide so-called trace formulae for determinations of observables to $\mathcal{O}(1/m_b^5)$ for $B \rightarrow X_c \ell \bar{\nu}$ Mannel, ISM, Vos [2311.12002]
- We built a model for QHDV guided by known contributions to $1/m_b^5$
Mannel, ISM, Verkade, Vos [soon!]
 - We model the factorial growth of the number of HQE parameters
 - We make an educated ansatz for the vQ dependence of $T_{\mu\nu}$ based on known contributions
 - We define the QHDV using the ambiguity arising from the inverse Borel transform
- Work in progress:
 - Full $|V_{cb}|^{\text{incl}}$ fit to $1/m_b^5$
 - Effect of QHDV on kinematic moments
 - Constructing observables sensitive to QHDV → test our model on data

Back-up 1: Cancellation without LLSA

- Assume for all dimension 8 operators that $X_i^5 \sim \Lambda_{\text{QCD}}^5$ and vary signs:



Back-up 2: Calculation of forward matrix element

- **Step 1:** expand charm propagator ($Q^\mu = m_b v^\mu$)

$$\begin{aligned}-iS_{\text{BGF}} &= \frac{1}{Q + iD - m_c} \\&= \frac{1}{Q - m_c} - \frac{1}{Q - m_c}(iD) \frac{1}{Q - m_c} \\&\quad + \frac{1}{Q - m_c}(iD) \frac{1}{Q - m_c}(iD) \frac{1}{Q - m_c} + \dots\end{aligned}$$

Back-up 2: Calculation of forward matrix element

- **Step 2:** insert in forward matrix element

$$\begin{aligned} T = & \left[\Gamma \frac{1}{\not{Q} - m_c} \Gamma^\dagger \right]_{\alpha\beta} \langle \bar{b}_\alpha b_\beta \rangle \\ & - \left[\Gamma \frac{1}{\not{Q} - m_c} \gamma^\mu \frac{1}{\not{Q} - m_c} \Gamma^\dagger \right]_{\alpha\beta} \langle \bar{b}_\alpha (iD_\mu) b_\beta \rangle \\ & + \left[\Gamma \frac{1}{\not{Q} - m_c} \gamma^\mu \frac{1}{\not{Q} - m_c} \gamma^\nu \frac{1}{\not{Q} - m_c} \Gamma^\dagger \right]_{\alpha\beta} \langle \bar{b}_\alpha (iD_\mu) (iD_\nu) b_\beta \rangle \\ & + \dots \end{aligned}$$

Back-up 2: Calculation of forward matrix element

- **Step 3:** Determine **Trace formula**

- Start at dimension 8 → lengthy, but systematically calculable
- Compute dim-7, including $1/m_b$ correction through e.o.m.

$$(ivD)b_v = -\frac{1}{2m_b}(i\not{D})(i\not{D})b_v$$

- ...
- Compute dim-3, including corrections up to $1/m_b^5$

$$\langle \bar{b}_\alpha b_\beta \rangle = 2m_B \left(\frac{1+\gamma}{4} + \frac{1}{8m_b^2} (\mu_G^2 - \mu_\pi^2) + \mathcal{O}(1/m_b^6) \right)_{\beta\alpha}$$

- Need full (non-RPI) set of **basic parameters** up to $1/m_b^5$

- **Step 4:** Compute the trace with the geometric series

Back-up 3:LLSA

- At $1/m_b^5 \rightarrow$ too many HQE parameters to fit to data
- **LLSA:** Lowest-Lying-State Approximation to estimate through known parameters μ_π^2 , μ_G^2 , ρ_D^3 and ρ_{LS}^3

Mannel, Turczyk, Uraltsev [1009.4622]

$$\langle B | \bar{b} \ A C \Gamma \ b(0) | B \rangle = \frac{1}{2m_B} \sum_n \langle B | \bar{b} \ A \ b(0) | n \rangle \cdot \langle n | \bar{b} \ C \Gamma \ b(0) | B \rangle$$

- $A = iD_{\mu_1} \dots iD_{\mu_k}$, $C = iD_{\mu_{k+1}} \dots iD_{\mu_n}$

Back-up 4: QHDV model

$$-\frac{1}{\pi} \hat{\Delta}_{DV} \text{Im} [T_{2,3}(vQ, Q^2)] =$$



$$C_{DV} \frac{N}{\Lambda_{HQE} - vQ} \frac{\Lambda_{HQE}}{\sqrt{Q^2}} \left(\sin \left(\frac{\sqrt{Q^2}}{\Lambda_{HQE}} \right) - \sqrt{\frac{vQ}{\Lambda_{HQE}}} \sin \left(\frac{1}{\sqrt{\Lambda_{HQE}}} \sqrt{\frac{Q^2}{vQ}} \right) \right)$$

$$-\frac{1}{\pi} \hat{\Delta}_{DV} \text{Im} [T_5(vQ, Q^2)] =$$

$$C_{DV} \frac{N}{\Lambda_{HQE} - vQ} \frac{(vQ)^2}{\Lambda_{HQE} \sqrt{Q^2}} \left(\sin \left(\frac{\sqrt{Q^2}}{\Lambda_{HQE}} \right) - \sqrt{\frac{\Lambda_{HQE}}{vQ}} \sin \left(\frac{1}{\sqrt{\Lambda_{HQE}}} \sqrt{\frac{Q^2}{vQ}} \right) \right)$$



Back-up 5: QHDV in q^2 moments

- Preliminary



Back-up 5: QHDV in q^2 moments

- Preliminary



Back-up 6: Dependence on Λ_{HQE}

- Preliminary

