


Lattice meets Continuum

Monday 30 September 2024 - Thursday 3 October 2024



Book of Abstracts

Contents

Short Flow-Time Expansion of the LEFT basis: Background Field Method and Chiral Symmetry	1
Continuum approaches to semileptonic $s \rightarrow d$ and $b \rightarrow s(d)$ transitions	1
Gradient Flow Renormalisation for Meson Mixing and Lifetimes	1
Semileptonic B decays into final states with heavy sterile neutrinos	2
Radiative corrections to $B \rightarrow \ell\nu$	2
Sum Rules for Lifetimes	2
Extracting excited-state contributions of the B_s to D_s semi-leptonic decays from inclusive lattice simulations	3
Factorial growth of perturbation theory and inclusive semileptonic B decays	3
Form factors for semi-leptonic $B_{(s)} \rightarrow D_{(s)}^* \ell\nu_\ell$ decays	4
Inclusive charm and tau decays	4
Semileptonic Inclusive Decay of the D_s Meson from Lattice QCD	4
Semileptonic B-> D^* decays	5
Semileptonic B-> D^* decays	5
Semileptonic B-> D^* decays	5
Semileptonic decays in the continuum	5
Vcb using inclusive q^2 moments	5
Inclusive decays on the lattice	6
Neutral meson mixing from lattice QCD	6
Meson mixing and lifetimes	6
Hadron spectroscopy and exotic states from lattice QCD	6
Hadron spectroscopy and exotic states in the continuum	6
Decays with 2 or more hadronic final states	6

Gradient flow and the short flow time expansion	6
Matching perturbation theory and lattice	7
Light cone sum rules and lattice QCD	7
Experimental overview: $ V_{cb} $ & $ V_{ub} $ at Belle and Belle II	7
Excited $B^*\pi$ states in $B \rightarrow \pi$ decays	7
Spectroscopy of tetra-quarks and other exotic states	7
Semileptonic decays lattice QCD	7
HQET on the lattice	7
Welcome	8
TBA	8
CKM unitarity and a truly global EFT analysis	8
TBA	8
TBA	8
A look into the $f_0(980)$ through the lens of rare B meson decays	8

Afternoon 2 / 1

Short Flow-Time Expansion of the LEFT basis: Background Field Method and Chiral Symmetry**Author:** Óscar Lara Crosas¹**Co-author:** Peter Stoffer¹¹ *UZH and PSI***Corresponding Authors:** oscar.laracrosas@physik.uzh.ch, stoffer@physik.uzh.ch

In this talk, I will present the Short Flow-Time expansion (SFTE) of operators present in the Low Energy Effective Field Theory (LEFT) up to dimension six, in the HV scheme. To be able to perform an off-shell calculation and avoid mixing with gauge-variant operators, we make use of the Background Field Method applied to the Gradient Flow. Moreover, we restore chiral symmetry by introducing appropriate counterterms. This work paves the way toward systematic higher-order SFTEs of LEFT operators, a key aspect to connect lattice simulations to continuum studies, which is a necessary ingredient e.g. for a robust determination of new-physics contributions to the neutron electric dipole moment.

Afternoon 2 / 2

Continuum approaches to semileptonic $s \rightarrow d$ and $b \rightarrow s(d)$ transitions**Author:** Jack Jenkins¹¹ *University of Siegen***Corresponding Author:** jack.jenkins@uni-siegen.de

In this talk I will discuss a continuum approach to the nonlocal contribution to rare kaon decays such as $K^+ \rightarrow \pi^+ \nu \nu$, with an emphasis on whether and how to decouple the charm. I will also motivate determinations of isospin breaking effects in $\Delta B = 0$ four-quark operators as input for theoretical predictions of inclusive semileptonic $b \rightarrow s(d)$ decays.

Afternoon 2 / 3

Gradient Flow Renormalisation for Meson Mixing and Lifetimes**Authors:** Andrea Shindler¹; Antonio Rago²; Fabian Lange³; Jonas Kohnen¹; Matthew Black⁴; Oliver Witzel⁵; Robert Harlander¹¹ *RWTH Aachen University*² *University of Southern Denmark*³ *Universität Zürich and Paul Scherrer Institut*⁴ *University Siegen*⁵ *Universität Siegen***Corresponding Authors:** shindler@physik.rwth-aachen.de, jonas.kohnen@rwth-aachen.de, oliver.witzel@uni-siegen.de, fabian.lange@physik.uzh.ch, rago@sdu.dk, robert.harlander@rwth-aachen.de, matthew.black@uni-siegen.de

The gradient flow offers a unique renormalization scheme applicable in both perturbation theory and lattice calculations. Especially, it suppresses operator mixing on the lattice, shifting it to the perturbative matching calculation. In this talk, we discuss using the gradient flow scheme for the four-quark operators describing neutral meson mixing and lifetimes. While meson mixing calculations are well-established on the lattice and validate our procedure, a lattice calculation of matrix elements for heavy meson lifetimes remains outstanding. We provide an overview of the full calculation process and present some preliminary results.

Afternoon 1 / 4

Semileptonic B decays into final states with heavy sterile neutrinos

Authors: Florian Bernlochner^{None}; Marco Fedele^{None}; Markus Prim^{None}; Tim Kretz^{None}; Ulrich Nierste¹

¹ *KIT*

Corresponding Author: ulrich.nierste@kit.edu

$B \rightarrow D^* \ell \nu$ decays are sensitive to contributions in which the missing energy and momentum stem from a hypothetical heavy sterile neutrino N . Belle II data on angular distributions in this decay are used to search for hints $B \rightarrow D^* \ell N$ in a model-independent way. To this end dimension-6 operators with different Dirac structures are considered and competitive upper bounds on some of their couplings are derived. I will show the allowed regions in the coupling-versus-mass plane and discuss the implications.

Afternoon 1 / 5

Radiative corrections to $B \rightarrow \ell \nu$

Author: Max Ferré¹

¹ *JGU (Mainz)*

Corresponding Author: ferremax@uni-mainz.de

In this talk I will focus on the study of the leptonic $B \rightarrow \ell \nu$ decay at next-to-leading order in QED. The future improvements of experimental measurements of this channel require a reliable theory prediction, hence a careful theoretical estimate of QED corrections. The multi-scale character of this process requires an appropriate effective theory (EFT) construction to factorize the different contributions. In the first part of this talk, I will discuss the EFT description of the process at the partonic level, which is based on Heavy Quark Effective Theory and Soft Collinear Effective Theory. I will show how the inclusion of QED corrections demands a generalisation of the hadronic decay constant defining a new non-perturbative input. In the second part of the talk, I will discuss the EFT description below the confinement scale based on the Chiral Lagrangian including Heavy Mesons (B and B^*). I will show that depending on the cut on final state radiation and on the lepton flavor the contribution from excited states of the B meson can become important.

Afternoon 1 / 6

Sum Rules for Lifetimes

Authors: Alexander Lenz¹; Martin Lang²; Matthew Black³; Zachary Wüthrich⁴

¹ *Siegen University*

² *TP1*

³ *University Siegen*

⁴ *Siegen*

Corresponding Authors: alexander.lenz@uni-siegen.de, martin.lang@uni-siegen.de, matthew.black@uni-siegen.de, zachary.wuethrich@uni-siegen.de

Precise determination of hadronic matrix elements plays a crucial role for interpreting potential deviations from the Standard Model observed in experiments testing flavor physics. While lattice QCD provides first principles calculations, current results are still limited to a subset of the operators that may appear in theories of new physics. The sum rule approach allows for a complementary determination of matrix elements directly from QCD, with theoretical uncertainties that can be systematically improved. Previous research successfully ascertained Standard Model hadronic matrix elements for dimension-six $F=0,2$ operators, demonstrating competitiveness with lattice findings. Our aim is to expand upon these findings by including the entire set of four-quark QCD operators for lifetimes, crucial in scenarios Beyond the Standard Model, where lattice results are currently absent. This extension includes operators with Dirac structures not previously examined in sum rules analyses documented in existing literature. This will provide for the first time bag parameter results which can increase the precision of a wide variety of new physics theories. The bag parameter results will be determined using HQET sum rules for three-point correlators, which requires a three-loop computation. In addition there is a one-loop computation of the QCD-HQET matching required.

Afternoon 1 / 7

Extracting excited-state contributions of the B_s to D_s semi-leptonic decays from inclusive lattice simulations

Authors: Adreas Juettner¹; Ahmed Elgaziari¹; Alessandro Barone²; Ryan Kellermann³; Shoji Hashimoto³; Takashi Kaneko³; Zhi HU³

¹ *University of Southampton*

² *Johannes Gutenberg University Mainz*

³ *High Energy Accelerator Research Organization (KEK)*

Corresponding Author: huzhi@post.kek.jp

We present a pilot study on extracting the P-wave form factors of the B_s to D_s semi-leptonic decays from the B_s four-point correlators. With their inclusive nature, four-point correlators include all the exclusive states from B_s with valence content of c and s quarks. We access the excited-state contributions by carefully fitting the correlators using multiple exponentials. In this pilot study, 2+1-flavour domain-wall fermion actions with approximately physical masses are utilized for light quarks. Heavy quarks, c and b , are simulated using relativistic-heavy quark actions. A single lattice ensemble with a lattice spacing of 0.11 fm is used for the analysis.

I am very eager to participate in this event, but currently, I am still seeking funding for the trip if my abstract gets accepted.

Afternoon 1 / 8

Factorial growth of perturbation theory and inclusive semileptonic B decays

Author: Andreas Kronfeld¹

¹ *Fermilab***Corresponding Author:** ask@fnal.gov

TBA

Afternoon 1 / 9**Form factors for semi-leptonic $B_{(s)} \rightarrow D_{(s)}^* \ell \nu_\ell$ decays****Author:** Anastasia Boushmelev¹**Co-authors:** Matthew Black²; Oliver Witzel³¹ *University of Siegen*² *University Siegen*³ *Universität Siegen***Corresponding Authors:** anastasia.boushmelev@uni-siegen.de, oliver.witzel@uni-siegen.de, matthew.black@uni-siegen.de

Semileptonic $B_{(s)}$ decays are of great phenomenological interest because they allow to extract CKM matrix elements or test lepton flavor universality. Taking advantage of existing data, we explore extracting form factors for vector final states using the narrow width approximation. Based on RBC-UKQCD's set of 2+1 flavor gauge field ensembles with Shamir domain-wall fermion and Iwasaki gauge field action, we study semileptonic $B_{(s)}$ decays using unitary light and strange quarks, Möbius domain wall fermions for charm quarks, and bottom quarks simulated with the relativistic heavy quark (RHQ) action. Exploratory results for $B_s \rightarrow D_s^* \ell \nu_\ell$ are presented.

Morning 2 / 10**Inclusive charm and tau decays****Author:** Alessandro De Santis¹¹ *University of Roma Tor Vergata and INFN***Corresponding Author:** alessandro.desantis@roma2.infn.it

The computation of observables from inclusive hadronic processes is widely regarded as a challenging task from the lattice QCD perspective. However, the access to such observables at the non-perturbative level is crucial, especially in Flavour Physics, where high-precision comparisons of first-principles theoretical predictions and experimental measurements could reveal discrepancies, potentially pointing to New Physics Scenarios beyond the Standard Model. A viable practical solution is provided by the Hansen-Lupo-Tantalo reconstruction method from Euclidean lattice correlation functions. Two recent applications of this method include the hadronic decays of the τ lepton, allowing for the extraction of the V_{us} and V_{ud} CKM matrix elements, and the semi-leptonic decay of the charmed D_s meson. Alongside a discussion of the methodology, we illustrate results for both the applications.

Afternoon 2 / 11**Semileptonic Inclusive Decay of the D_s Meson from Lattice QCD**

Authors: Alessandro De Santis¹; Christiane Groß²

¹ *University of Roma Tor Vergata and INFN*

² *HISKP, Uni Bonn*

Corresponding Authors: gross@hiskp.uni-bonn.de, alessandro.desantis@roma2.infn.it

We analyse the decay of the D_s meson, focusing on the $D_s \rightarrow X\ell\nu$ channel based on Extended Twisted mass ensembles at the physical pion mass value.

On the basis of four-point correlation functions, we use the HLT reconstruction method to calculate the differential decay rate and differential first lepton energy moment.

The analysis is performed at four different lattice spacings, and three different volumes to address the associated systematic uncertainties.

We focus on the systematic errors arising from the procedure and include data for the first lepton energy moment.

Morning 2 / 12

Semileptonic B- \rightarrow D* decays

Corresponding Author: alexv@unizar.es

Morning 2 / 13

Semileptonic B- \rightarrow D* decays

Corresponding Author: takashi.kaneko@kek.jp

Afternoon 1 / 14

Semileptonic B- \rightarrow D* decays

Corresponding Author: martin.jung@unito.it

Morning 1 / 15

Semileptonic decays in the continuum

Corresponding Author: florian.s.herren@gmail.com

Morning 1 / 16

Vcb using inclusive q^2 moments

Corresponding Author: kerivos@gmail.com

Morning 2 / 17

Inclusive decays on the lattice

Morning 2 / 18

Neutral meson mixing from lattice QCD

Corresponding Author: felix.erben@ed.ac.uk

Morning 2 / 19

Meson mixing and lifetimes

Corresponding Author: maria.piscopo@uni-siegen.de

Morning 1 / 20

Hadron spectroscopy and exotic states from lattice QCD

Corresponding Author: d.mohler@gsi.de

Morning 1 / 21

Hadron spectroscopy and exotic states in the continuum

Corresponding Author: c.hanhart@fz-juelich.de

Morning 1 / 22

Decays with 2 or more hadronic final states

Corresponding Author: fernando.romero-lopez@unibe.ch

Afternoon 1 / 23

Gradient flow and the short flow time expansion

Corresponding Author: robert.harlander@rwth-aachen.de

Afternoon 1 / 24

Matching perturbation theory and lattice

Corresponding Author: martin.gorbahn@liverpool.ac.uk

Morning 1 / 25

Light cone sum rules and lattice QCD

Corresponding Author: khodjamirian@physik.uni-siegen.de

Morning 1 / 26

Experimental overview: $|V_{cb}|$ & $|V_{ub}|$ at Belle and Belle II

Corresponding Author: raynette.vantonder@mcgill.ca

Morning 2 / 27

Excited $B^* \pi$ states in $B \rightarrow \pi$ decays

Corresponding Author: obaer@physik.hu-berlin.de

Morning 2 / 28

Spectroscopy of tetra-quarks and other exotic states

Corresponding Author: vadim.baru@tp2.rub.de

Morning 1 / 29

Semileptonic decays lattice QCD

Corresponding Author: j.t.tsang@cern.ch

Afternoon 2 / 30

HQET on the lattice

Corresponding Author: heitger@uni-muenster.de

Morning 1 / 31

Welcome

Corresponding Authors: oliver.witzel@uni-siegen.de, mannel@physik.uni-siegen.de

Afternoon 2 / 32

TBA

Afternoon 2 / 33

CKM unitarity and a truly global EFT analysis

Corresponding Author: tiantian.tong@uni-siegen.de

Afternoon 2 / 35

TBA

Afternoon 2 / 36

TBA

Afternoon 2 / 37

A look into the $f_0(980)$ through the lens of rare B meson decays

Author: Jaime del Palacio Lirola¹

¹ *University of Siegen*

Corresponding Author: jaime.delpalaciolirola@uni-siegen.de

The nature of many experimentally observed hadronic resonances has been a topic of debate ever since the 1960's. In this work, we focus on studying the $f_0(980)$, the second lightest unflavoured scalar resonance. We comment on the different descriptions for it in the quark model, and focus specifically on the pure $s\bar{s}$ picture. We mainly explore the $B_s^0 \rightarrow f_0(980)\mu^+\mu^-$ decay. We analyse the impact of the form factors, which are difficult to determine theoretically. They come from the hadronic matrix element and contain the hadronic information of the decay. Using the framework of the Weak Effective Theory, we utilize Wilson coefficients derived from observables of rare decays with the same quark-level transition, like $B \rightarrow K^{(*)}\mu^+\mu^-$, to probe the hadronic nature of the $f_0(980)$, even in the presence of possible New Physics. The effect of different theoretical form factor calculations on several observables is explored in detail. A range for the experimental untagged

branching ratio integrated in the $[1, 6] q^2$ bin (where q^2 is the square of the 4-momenta of the muon pair) is found to be $\mathcal{BR}_{\text{exp}} \in [0.04, 2.11] \times 10^{-7}$, according to the current form factor calculations, which all assume a pure $s\bar{s}$ state. The viability of extracting the form factors from experimental data is also studied, finding that determining one of them from the branching ratio would be possible with good precision.