



GW Transient Catalog 4.0 by the LVK collaboration



218 compact-binary mergers, data from observing runs O1, O2, O3, O4a

.


GW Transient Catalog 4.0 by the LVK collaboration




218 compact-binary mergers, data from observing runs O1, O2, O3, O4a

Pulsar timing arrays (PTAs)

Array of pulsars across our MW spiral arm \rightarrow GW detector of galactic dimensions!





[nrao.edu]

[MPIfR]

Pulsars: Highly magnetized rotating neutron stars

- \blacksquare Beamed radio pulses emitted from magnetic N and S poles \to cosmic lighthouses
- lacksquare Stable rotation with periods as short as a few milliseconds ightarrow celestial clocks

Look for tiny distortions in pulse times of arrival (TOAs) caused by nanohertz GWs

,

2023 PTA results



EPTA: European PTA
CPTA: Chinese PTA
PPTA: Parkes PTA
InPTA: Indian PTA
MPTA: MeerKAT PTA
NANOGrav: North American
Nanohertz Observatory for
Gravitational Waves


18 papers on the arXiv on June 29, 2023

[2306.16213]	NANOGrav	GWB	[2306.16222]	NANOGrav	Continuous GW
	IVAIVOGIAV	GVVD		IVAIVOGIAV	Continuous GVV
[2306.16214]	EPTA	GWB	[2306.16223]	NANOGrav	Analysis pipeline
[2306.16215]	PPTA	GWB	[2306.16224]	EPTA	Data set
[2306.16216]	CPTA	GWB	[2306.16225]	EPTA	Noise model
[2306.16217]	NANOGrav	Data set	[2306.16226]	EPTA	Continuous GW
[2306.16218]	NANOGrav	Noise model	[2306.16227]	EPTA	Implications
[2306.16219]	NANOGrav	New physics	[2306.16228]	EPTA	ULDM
[2306.16220]	NANOGrav	SMBHBs	[2306.16229]	PPTA	Noise model
[2306.16221]	NANOGrav	Anisotropies	[2306.16230]	PPTA	Data set

4


Evidence for Hellings–Downs (HD) correlations $\Gamma(\xi_{ab})$

2306.16213: NANOGrav




68 pulsars, 16 yr of data, HD at \sim 3 \cdots 4 σ

2306.16215: PPTA




32 pulsars, 18 yr of data, HD at \sim 2 σ

2306.16214: EPTA+InPTA



25 pulsars, 25 yr of data, HD at \sim 3 σ


2306.16216: CPTA



57 pulsars, 3.5 yr of data, HD at \sim 4.6 σ


Common power spectrum $\sqrt{\Delta f P_g(f)}$

2306.16213: NANOGrav




68 pulsars, 16 yr of data, HD at \sim 3 \cdots 4 σ

2306.16215: PPTA



32 pulsars, 18 yr of data, HD at \sim 2 σ

2306.16214: EPTA+InPTA



25 pulsars, 25 yr of data, HD at \sim 3 σ


2306.16216: CPTA



57 pulsars, 3.5 yr of data, HD at \sim 4.6 σ


Power-law parameters ${\it A}$ and γ

2306.16213: NANOGrav




68 pulsars, 16 yr of data, HD at \sim 3 \cdots 4 σ

2306.16215: PPTA




32 pulsars, 18 yr of data, HD at \sim 2 σ

2306.16214: EPTA+InPTA




25 pulsars, 25 yr of data, HD at \sim 3 σ

2306.16216: CPTA



57 pulsars, 3.5 yr of data, HD at \sim 4.6 σ

IPTA comparison





- Results from regional PTAs are consistent with each other (1σ posteriors overlap)
- Joint posterior = naive product (properly normalized) of individual posteriors
- lacktriangle Proper data combination and combined data analysis ightarrow IPTA DR3

8

MeerKAT Pulsar Timing Array (MPTA): 83 pulsars, 4.5 yr of data

[MPTA 2412.01153]






Abstract: [...] Under different assumptions about noise processes we can produce either what appear to be compelling Hellings-Downs correlations of high significance $(3-3.3\,\sigma)$ with a spectrum close to that which is predicted, or surprisingly, under slightly different assumptions, ones that are insignificant. This appears to be related to the fact that many of the highest precision MeerKAT Pulsar Timing Array pulsars are in close proximity and dominate the detection statistics [...]

Interpretations

• Supermassive black-hole binaries



2 GWs from the Big Bang



• SMBHBs (realistic)

- \blacksquare No SMBHB mergers directly observed as of yet \to data-driven field thanks to PTAs

New physics (speculative)


- Logical possibility: PTA signal is not of SMBHB origin or receives several contributions
- Probe and constrain cosmology at early times as well as particle physics at high energies

BSM scenario: Cosmic inflation

Big questions: What set the initial conditions of the Hot Big Bang: homogeneity, isotropy, spatial flatness? What seeded the temperature fluctuations in the CMB?

1 BSM scenario: Cosmic inflation

Big questions: What set the initial conditions of the Hot Big Bang: homogeneity, isotropy, spatial flatness? What seeded the temperature fluctuations in the CMB?



Cosmic inflation: Stage of exponentially fast expansion before the Hot Big Bang


- Requires form of dark energy, e.g., potential energy of a scalar "inflaton" field
- ullet Inflaton and metric fluctuations o primordial scalar and tensor perturbations

Primordial tensor spectrum

$$\mathcal{P}_t = r A_s \left(\frac{f}{f_{\mathrm{cmb}}}\right)^{n_t}$$

Parameters

 $T_{\rm rh}$ Reheating temperature r Tensor-to-scalar ratio n_t Tensor spectral index




Lessons

- Strongly blue-tilted spectrum, $n_t \sim 2 \cdots 4 \rightarrow \text{probe nonminimal inflation models}$
- ullet Transition from **reheating** to the Hot Big Bang in the PTA band for $T_{
 m rh}\sim 1\,{
 m GeV}$
- If GWB extrapolated to higher frequencies ightarrow large contribution to dark radiation

Example: Axion spectator dynamics during inflation

[Obata, Papagerogiou, Ünal 2307.02322]




Inflaton ϕ + axion spectator χ + Abelian gauge field A_{μ}

$$\mathcal{L} = \frac{1}{2} \partial_{\mu} \phi \, \partial^{\mu} \phi + \frac{1}{2} \partial_{\mu} \chi \, \partial^{\mu} \chi - \frac{1}{4} \frac{\mathsf{F}_{\mu\nu} \mathsf{F}^{\mu\nu}}{\mathsf{F}^{\mu\nu}} - V(\phi) - V(\chi) - \lambda \frac{\chi}{f} \frac{1}{4} \frac{\mathsf{F}_{\mu\nu} \tilde{\mathsf{F}}^{\mu\nu}}{\mathsf{F}^{\mu\nu}}$$

- Axion spectator field rolls down potential $V(\chi) \propto \cos(\chi/f)$ at some point during inflation
- Gauge-field modes with either positive or negative helicity become tachyonically unstable
- GWs from vacuum tensor perturbations + classical electric and magnetic fields

GWs from axion inflation

[von Eckardstein, KS, Sobol 2508.00798] [von Eckardstein, KS, Sobol 2509.25013] [von Eckardstein 2510.12644] New software package: Gradient Expansion Formalism Factory (GEFF)



Axion inflaton χ + Abelian gauge field A_{μ} (+ charged fermions ψ_{i})

$$\mathcal{L} = \frac{1}{2} \partial_{\mu} \chi \, \partial^{\mu} \chi - \frac{1}{4} \frac{\mathbf{F}_{\mu\nu} \mathbf{F}^{\mu\nu}}{\mathbf{F}^{\mu\nu}} - V\left(\chi\right) - \lambda \frac{\chi}{f} \frac{1}{4} \frac{\mathbf{F}_{\mu\nu} \tilde{\mathbf{F}}^{\mu\nu}}{\mathbf{F}^{\mu\nu}} - \mathbf{A}_{\mu} \mathbf{J}_{\psi}^{\mu}$$

- Pure axion inflation (PAI): GW production at the end of inflation; too much dark radiation
- Fermionic axion inflation (FAI): LISA/CE/ET target; PTA explanation in conflict with CMB

BSM scenario: Primordial black holes

Big questions: Are some of the black holes seen by LVK of primordial origin? To what extent do PBHs contribute to dark matter? How do galactic SMBHs form?

2 BSM scenario: Primordial black holes

Big questions: Are some of the black holes seen by LVK of primordial origin? To what extent do PBHs contribute to dark matter? How do galactic SMBHs form?



PBHs: Form in the gravitational collapse of large overdensities in the early Universe


- Typical scenario: Scalar perturbations enhanced during ultra-slow-roll inflation
- ullet Enhanced scalar perturbations o GWs at second order in perturbation theory

Primordial scalar spectrum

$$\mathcal{P}_{s} = rac{A}{\sqrt{2\pi}\,\Delta}\,\exp\left[-rac{(\ln{(f/f_*)})^2}{2\,\Delta^2}
ight]$$



- Peak frequency
- Peak amplitude
- Width



Lessons


- Require large-amplitude peak in $\mathcal{P}_s \to \text{input for building models of inflation}$
- PBH dark matter might be possible; but some tension with PBH overproduction
- On-going debate on impact of non-Gaussianities on efficiency of PBH production

BSM scenario: Phase transition

Big questions: How are the Higgs mechanism and the quark—hadron transition realized in the early Universe? Are there other fundamental forces beyond the Standard Model?

3 BSM scenario: Phase transition

Big questions: How are the Higgs mechanism and the quark—hadron transition realized in the early Universe? Are there other fundamental forces beyond the Standard Model?



Cosmological phase transitions: Changes in the quantum field theory vacuum structure

- SM predicts smooth crossovers; strong first-order phase transitions require BSM
- GWs from bubble collisions, sound waves, and magnetohydrodynamic turbulence

Peak amplitude and frequency


$$\begin{split} &\Omega_{\rm GW}^{\rm peak} \propto (H_* R_*)^2 \left(\frac{\alpha_*}{1+\alpha_*}\right)^2 \\ &f_{\rm peak} \propto \frac{T_*}{H_* R_*} \end{split}$$

Parameters

 T_* Percolation temperature

 α_* Transition strength

 H_*R_* Bubble separation



Lessons

- Strong $(\alpha_* \sim 1)$ and slow $(H_*R_* \sim 1)$ transition at a temperature $T_* \sim 100\,\mathrm{MeV}$
- Just the right ballpark for BSM modifications of the QCD phase transition
- lacktriangledown Alternatively, phase transition in a dark sector ightarrow complementary to lab searches

BSM scenario: Cosmic defects

Big questions: How are the tiny SM neutrino masses generated? What is the origin of the matter—antimatter asymmetry? Is the SM embedded in a grand unified theory?

4 BSM scenario: Cosmic defects

Big questions: How are the tiny SM neutrino masses generated? What is the origin of the matter–antimatter asymmetry? Is the SM embedded in a grand unified theory?



Cosmic strings / domain walls: Defects after spontaneous breaking of GUT symmetries

- ullet Typical scenario: $U(1)_{B-L}$ breaking o neutrino masses, leptogenesis, and strings
- Dynamics and decay of defect networks yield anisotropic stress and hence GWs


Decay rate per length

$$\Gamma_d = \frac{\mu}{2\pi} e^{-\pi\kappa}$$

Parameters

Tension (energy per length)


 κ Decay parameter



Lessons


- Prefered parameter values ightarrow input for GUT model building at $E \lesssim 10^{16} \, {
 m GeV}$
- Metastable strings yield a good fit; can be probed / excluded by LVK observations
- PTA bounds outperform CMB bounds, irrespective of the origin of the signal (!)

Bayesian model comparison



- Many BSM models reach Bayes factors of the order of $10 \cdots 100$
- Interesting but not conclusive; lots of uncertainties in SMBHB and BSM models


Bayesian model comparison



- ullet Many BSM models reach Bayes factors of the order of $10\cdots 100$
- Interesting but not conclusive; lots of uncertainties in SMBHB and BSM models


Call to action: Improve modelling on both the astro and the cosmo side!


PTArcade



Our code developed for 2306.16219: Fit your favorite BSM model to the NG15 data! New functionalities, new models, and new data (when available) added on a steady basis

Complementary observables: Anisotropies







Search for anisotropies in the GWB signal in the sky

- Current sensitivity already at the level of expected anisotropies from SMBHBs
- No signal detected ightarrow sky-dependent upper limits on deviation from monopole

No detection of anisotropies with future data sets \rightarrow hint of primordial origin!?

Complementary observables: Continuous waves






Search for continuous-wave signals from individual nearby SMBHB systems

- Interesting hints in the data, which, however, do not withstand further scrutiny
- ullet Overall, no signal detected o sky-dependent upper limits on GW amplitude

This is only the beginning!



A bright future for GW science with PTAs

- \blacksquare Status: Common-spectrum process; $3\cdots 4\,\sigma$ evidence for HD correlations
- Next: HD correlations at 5σ , spectral shape, anisotropies across the sky, ...
- Promise: Deep insights into galaxy and BH evolution and/or new physics

Stay tuned!

And thanks a lot for your attention